

LOCAL COHOMOLOGIES OF ISOLATED NON F -RATIONAL SINGULARITIES

YUKIHIRO TAKAYAMA

ABSTRACT. In this paper, we consider positively graded isolated non F -rational singularities $(R, \mathfrak{m})$ with $d = \dim R$ over the field K of positive characteristic. We give a representation of lower local cohomologies $H_{\mathfrak{m}}^i(R)$ ($i < d$) in terms of tight closure and limit closure of certain type of parameters. As an application to isolated singularities, we show a relation between non-vanishing of the tight closure of zero in the highest local cohomology $(0)_{H_{\mathfrak{m}}^d(R)}^*$ and non-vanishing of the cohomology $H_{\mathfrak{m}}^{d-1}(R)$.

MSC Primary: 13A35, 13D45, Secondary: 13A02

INTRODUCTION

Let K be a field of characteristic $\text{char } K = p > 0$. Let $(R, \mathfrak{m})$ be a finitely generated $\mathbb{N}$ -graded K -algebra with $d = \dim R$, $R = \bigoplus_{n \geq 0} R_n$, $\mathfrak{m} = \bigoplus_{n > 0} R_n$ and $R_0 = K$. We assume that R is reduced, equidimensional and R_P is F -rational for all primes $P (\neq \mathfrak{m})$, namely R is an $\mathbb{N}$ -graded isolated non F -rational singularity. Since R is Cohen-Macaulay on the punctured spectrum, it is a generalized Cohen-Macaulay ring, i.e., $\ell_R(H_{\mathfrak{m}}^i(R)) < \infty$ for all $i < d$, where $H_{\mathfrak{m}}^i(R)$ is the i th local cohomology module and $\ell_R(-)$ denotes the length as a R -module. In this paper, we are interested in the structure of local cohomologies $H_{\mathfrak{m}}^i(R)$.

A distinguished property of $\mathbb{N}$ -graded isolated non F -rational singularity is Kodaira vanishing theorem, which, in terms of local cohomologies, can be stated that lower local cohomologies vanish for negative degrees [10, 11]: $[H_{\mathfrak{m}}^i(R)]_n = 0$ for all $i < d$ and all $n < 0$. If R is a generic mod p reduction of an isolated non-rational singularity (over the field of characteristic 0), Kodaira vanishing holds. This result has been established by Huneke, Smith, Hara and Watanabe [6, 11, 7]. This is in fact the case of *liftable to the second Witt vectors* in the sense of Deligne-Illusie [1]. Notice that an isolated non F -rational singularity is not necessary a mod p reduction of an isolated non-rational singularity and Kodaira vanishing for positive characteristic is in general false, eg. [15]. This means that vanishing/non-vanishing of local cohomologies for isolated non F -rational singularities in general is widely open.

In this paper, we first show that lower local cohomologies $H_{\mathfrak{m}}^i(R)$, $i < d$, of an $\mathbb{N}$ -graded isolated non F -rational singularity R can be described as quotient modules of the tight closures by the limit closures of suitable parameter ideals (see Theorem 17). This is a refinement of the results by Schenzel [12] and Smith [14]. With this representation, Huneke-Smiths interpretation of Kodaira vanishing in terms of tight closure [11] can be recovered immediately in a slightly general form.

A Noetherian local ring $(R, \mathfrak{m})$ is an isolated non F -rational singularity if and only if R is generalized Cohen-Macaulay and the tight closure of zero in the highest local cohomology $(0)_{H_{\mathfrak{m}}^d(R)}^*$, $d = \dim R$, has finite length (cf. [4]). This suggests that vanishing/non-vanishing of lower local cohomologies may be controlled by $(0)_{H_{\mathfrak{m}}^d(R)}^*$ to some extent. In fact, the above mentioned Kodaira vanishing theorem by Huneke-Smith-Hara-Watanabe has been proved by investigating the structure of $(0)_{H_{\mathfrak{m}}^d(R)}^*$. In this paper, we will consider isolated singularities and show that $[(0)_{H_{\mathfrak{m}}^d(R)}^*]^n \neq 0$ for some $n \in \mathbb{Z}$ implies non-vanishing of $H_{\mathfrak{m}}^{d-1}(R)$ for certain degree, under some subtle conditions (see Theorem 24).

In section 1 we will summarize the known results on isolated non F -rational singularities that are necessary in this paper. In section 2, after preparing some definitions and facts on standard sequence, limit closure and germ closure, we give a representation of lower local cohomologies in terms of tight closure and limit closure. We also give some direct consequences from this representation. $\mathbb{N}$ -graded isolated singularities are considered in section 3. After giving a graded version of Goto-Nakamura's representation of tight closure of zero $(0)_{H_{\mathfrak{m}}^d(R)}^*$, we investigate non-vanishing of $(0)_{H_{\mathfrak{m}}^d(R)}^*$ and a generic hypersurface intersection $\overline{R}$ of R . Here we use the characteristic p version of Flenner's Bertini theorem. Then finally we consider the non-vanishing of $H_{\mathfrak{m}}^{d-1}(R)$ in terms of non-vanishing of $(0)_{H_{\mathfrak{m}}^d(R)}^*$.

1. GRADED ISOLATED NON F -RATIONAL SINGULARITIES

This section summarizes the basic definitions and results concerning isolated non F -rational singularities. See, for example, [10, 9, 14] for the detail.

Definition 1. A sequence $x_1, \dots, x_i$ in a ring R is called parameters if for every $P \in \operatorname{Spec} R$ containing the sequence, the image of the sequence in R_P forms a part of sop. In other words, $\operatorname{ht}((x_1, \dots, x_i)) = i$. An ideal generated by a set of parameters is called a parameter ideal.

Definition 2. For an ideal I of a ring R , the tight closure I^* of I is defined by $I^* = \{r \in R \mid \text{there exists } c \in R^\circ \text{ such that } cz^q \in I^{[q]} \text{ for all } q = p^e \gg 0\}$, where R° is the set of elements outside the union of minimal primes of R and $I^{[q]} = (r^q \mid r \in I)$.

Definition 3. A Noetherian ring R of prime characteristic is said to be F -rational if every parameter ideal of R is tightly closed.

We recall the basic properties of F -rational rings needed in this paper.

Proposition 1 (cf. [9]). Let R be a Noetherian ring that is the homomorphic image of a Cohen-Macaulay ring. We have the following:

- (1) If R is F -rational, then R is Cohen-Macaulay.
- (2) A localization of an F -rational ring is F -rational.
- (3) If R is regular, then R is F -rational.

Definition 4. Let K be a field. Then a finitely generated $\mathbb{N}$ -graded K -algebra $(R, \mathfrak{m})$ is said to be a generalized Cohen-Macaulay if one of the following equivalent conditions holds:

- (1) $\text{Proj}(R)$ is an equidimensional Cohen-Macaulay projective scheme,
- (2) R_P is an equidimensional Cohen-Macaulay local ring for every $P \in \text{Spec } R - \{\mathfrak{m}\}$,
- (3) $H_{\mathfrak{m}}^i(R)$ is a finite length module for all $i < \dim R$. In particular, there exists $N_i \in \mathbb{Z}$ such that $[H_{\mathfrak{m}}^i(R)]_n = 0$ for all $n < N_i$.

Definition 5. Let $(R, \mathfrak{m})$ be a finitely generated $\mathbb{N}$ -graded K -algebra or a Noetherian local ring with the maximal ideal $\mathfrak{m}$. Then we say that R is an isolated non F -rational singularity if R_P is F -rational for all primes $P (\neq \mathfrak{m})$.

In this paper, when we consider an isolated non F -rational singularity, we always assume that it is equidimensional.

Definition 6. An element $c \in R^\circ$ is called a parameter test element if, for arbitrary ideal I generated by an sop and arbitrary $x \in I^*$, we have $cx^q \in I^{[q]}$ for all $q = p^e$.

In this definition, we can replace the ideal I by any parameter ideal, including $I = (0)$, because of the following result, which is a special case of Exercise 2.12 [9].

Proposition 2. Let $(R, \mathfrak{m})$ be a local ring and assume that $c \in R^\circ$ is a test element for parameter ideals generated by sops. Then c is a test element for arbitrary parameter ideal $I = (x_1, \dots, x_i)$, $i < d = \dim R$, including the case of $i = 0$, i.e., $I = 0$.

Every isolated non F -rational singularity has an $\mathfrak{m}$ -primary parameter test ideal. Namely,

Proposition 3. Let $(R, \mathfrak{m})$ be a reduced isolated non- F -rational singularity of $\text{char } R = p > 0$. Then there exists an $\mathfrak{m}$ -primary parameter test ideal $J \subset R$, i.e., every element $a \in J$ is a test element for any parameter ideal in R .

Proof. This follows immediately from (3.9) [16]. See also Exer. 2.12 [9]. $\square$

By Prop. 1, we know that an isolated non F -rational singularity is generalized Cohen-Macaulay. Also, in view of Prop. 3, we will always consider reduced rings.

2. LOWER LOCAL COHOMOLOGIES OF ISOLATED NON F -RATIONAL SINGULARITIES

This section gives a representation of lower local cohomologies of isolated non F -rational singularities in terms of tight closure and limit closure of unconditioned strong d -sequences (USD-sequences), which is a refinement of the representation given by P. Schenzel and K. E. Smith.

2.1. equidimensional hull, limit closure and germ closure. In this subsection, we summarize some of the definitions and results needed to give our representation of lower local cohomologies.

Definition 7. A sequence of elements $x_1, \dots, x_n$ in a commutative ring is said to be a d -sequence if for every $0 \leq i \leq n - 1$ and $k > i$ we have

$$(x_1, \dots, x_i) : x_{i+1}x_k = (x_1, \dots, x_i) : x_k.$$

A sequence $x_1, \dots, x_n$ is said to be a strong d -sequence if $x_1^{m_1}, \dots, x_n^{m_n}$ is a d -sequence for every arbitrary $m_i \geq 1$, $i = 1, \dots, n$. Finally, a sequence $x_1, \dots, x_n$ is said to be a USD-sequence (unconditioned strong d -sequence) if every permutation of it is a strong d -sequence.

Definition 8. Let $(R, \mathfrak{m})$ be a Noetherian local ring with $d = \dim R$. A system of parameters $x_1, \dots, x_d$ is called standard if

$$(x_1, \dots, x_d)H_{\mathfrak{m}}^j(R/(x_1, \dots, x_{i-1})) = 0 \quad \text{for all } 0 \leq i + j \leq d, i \geq 1$$

where we set $(x_1, \dots, x_{i-1}) = 0$ if $i = 1$.

Notice that for a standard sop we have

$$(x_1, \dots, x_d)H_{\mathfrak{m}}^i(R) = 0 \quad \text{for all } i < d.$$

We also note that the empty sequence is trivially a USD and standard sequence. The notions of USD-sequence and standard sequence are actually equivalent:

Proposition 4 (cf. (3.8) [12]). For an sop $x_1, \dots, x_d$ in a Noetherian local ring $(R, \mathfrak{m})$, the following are equivalent:

- (i) $x_1, \dots, x_d$ is a standard system of parameters;
- (ii) $x_1, \dots, x_d$ is an unconditioned strong d -sequence (USD-sequence).

The USD or standad property of the sequences is preserved by hypersurface intersection. Namely,

Proposition 5 (cf. (3.2) [12]). Let $x_1, \dots, x_i$ be a standard sequence in a Noetherian local ring $(R, \mathfrak{m})$. Then the image of $x_1, \dots, x_{i-1}$ in $\overline{R} := R/x_i R$ is again a standard sequence of $\overline{R}$.

For the existence of USD-sequence, we have

Proposition 6 ((6.19) [5]). Let $(R, \mathfrak{m})$ be a generalized Cohen-Macaulay local ring. Then there exists $t \in \mathbb{N}$ such that any system of parameters in $\mathfrak{m}^t$ is a USD-sequence.

This implies that we can always obtain a USD-sequence by taking hight enough power of an sop:

Corollary 7. Let $(R, \mathfrak{m})$ be a generalized Cohen-Macaulay local ring and let $x_1, \dots, x_d$, $d = \dim R$, be any system of parameters. Then $x_1^n, \dots, x_d^n$ is a USD-sequence for $n \gg 0$.

Proposition 8 (cf. (5.11) [10]). Let $x_1, \dots, x_i \in R$ be parameters which are also parameter test elements. Then $x_1, \dots, x_i$ are a USD-sequence.

Definition 9. Let R be a Noetherian ring. For an ideal $I \subset R$, consider the minimal primary decomposition $I = \bigcap_i Q_i$. Then we define $I^{umn} = \bigcap Q_i$, where Q_i runs over the primary components such that $\dim R/Q_i = \dim R/\sqrt{Q_i} = \dim R/I$. We call I^{umn} the unmixed hull of I .

Definition 10. For a parameter ideal $I = (x_1, \dots, x_i)$ of a ring R , we define the limit closure of I as follows. If $i \geq 0$, we define

$$\begin{aligned} I^{\lim} &= \{z \in R \mid \mathbf{x}^{s-1}z \in (x_1^s, x_2^s, \dots, x_i^s) \text{ for some } s \in \mathbb{N}\} \\ &= \{z \in R \mid \mathbf{x}^{s-1}z \in (x_1^s, x_2^s, \dots, x_i^s) \text{ for all } s \gg 0\} \\ &= \bigcup_{s=1}^{\infty} (x_1^s, \dots, x_i^s) : \mathbf{x}^{s-1} \end{aligned}$$

where $\mathbf{x} = x_1 \cdots x_i$. For $i = 0$, we define $(0)^{\lim} = 0$.

Proposition 9 (cf. (2.5) [11]). Let $(R, \mathfrak{m})$ be a Noetherian local ring and $I = (x_1, \dots, x_i)$ ($i \geq 0$) be a parameter ideal. Then the following are equivalent:

- (i) $z \in I^{\lim}$
- (ii) an element $\eta = \left[\frac{z}{x_1 \cdots x_i} \right] \in H_I^i(R)$ is 0.

Proof. Clear from the Čech complex representation of local cohomology. $\square$

Proposition 10 ((5.4) [10]). Let R be equidimensional and the homomorphic image of a Cohen-Macaulay ring. If $I \subset R$ is a parameter ideal, then we have $I^{\lim} \subseteq I^*$.

Proposition 11 ((5.8) [10]). Let $(R, \mathfrak{m})$ be an equidimensional graded Noetherian ring over a field with an $\mathfrak{m}$ -primary parameter test ideal. Then for a parameter ideal $I \subset R$ such that $\text{ht } I < \dim R$, we have $I^{\text{um}n} = I^*$.

Corollary 12. Let $(R, \mathfrak{m})$ be an $\mathbb{N}$ -graded isolated non F -rational singularity and let $I = (x_1, \dots, x_i)$ be a parameter ideal such that $\text{ht}(I) < \dim R$. Then we have $I^{\text{un}m} = I^*$.

Proof. Immediate from Prop. 3 and Prop. 11. $\square$

Definition 11. Let $I = (x_1, \dots, x_i)$ be a parameter ideal. Then we define the germ closure of I as follows. For $i \geq 1$, we define

$$I^{\text{germ}} = \sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^*.$$

Also, we define $(0)^{\text{germ}} = 0$.

In the above definition, we note that $I^{\text{germ}} = (0)^*$ if $i = 1$ and we have $I \subset I^{\text{germ}}$ for $i \neq 1$. Also we always have $I^{\text{germ}} \subset I^*$.

The following proposition is a slight modification of Theorem 5.12 [10]. The only difference is that, for $I = (x_1, \dots, x_i)$ for $i \leq d$, only the case of $i = d$ is considered in [10]. But with almost the same proof we have the following extension.

Proposition 13 (cf. (5.12) [10]). Let $(R, \mathfrak{m})$ be an equidimensional local ring that is the homomorphic image of a Cohen-Macaulay local ring. Let $I = (x_1, \dots, x_i)$, where $0 \leq i \leq d$ and $x_1, \dots, x_d$ is an sop which are parameter test elements. Then we have $I^{\text{germ}} + I = I^{\lim}$, in particular if $i \neq 1$ we have $I^{\text{germ}} = I^{\lim}$.

Proof. We only show the case of $i = 1$. Other cases are similar to (5.12) [10]. Let $x \in R$ be a parameter test element. We have $(x)^{germ} = (0)_R^*$ by the definition. Let $z \in (x)^{lim}$ be arbitrary. Then $x^s z \in (x^{s+1})$ for some $s \in \mathbb{N} \cup \{0\}$, so that $x^s(z - rx) = 0$ for some $r \in R$. Assume that $z - rx \neq 0$. Since x is a parameter test element, which also implies to be a test element for (0) , we have $z - rx \in (0)^*$ so that $z \in (x) + (0)^* = (x) + (x)^{germ}$. Thus we have $(x)^{lim} \subset (x) + (x)^{germ}$. For the converse inclusion, $(x) \subset (x)^{lim}$ is immediate. Again, since x is a parameter test element, we have, for $z \in (0)_R^* = (x)^{germ}$, $xz^q = 0$ for any $q = p^e$. In particular, $xz = 0 \in (x^2)$, which implies $z \in (x)^{lim}$. Consequently, we have $(x)^{lim} \supset (x) + (x)^{germ}$. $\square$

2.2. representation of lower local cohomologies.

Proposition 14 (cf. (6.8) and (6.9) [14]). *Let $(R, \mathfrak{m})$ be a Noetherian local ring that is equidimensional, the homomorphic image of a Cohen-Macaulay local ring and R_P is Cohen-Macaulay for all primes $P \neq \mathfrak{m}$. Then there exists an sop $x_1, \dots, x_d$, ($d = \dim R$) such that for $0 \leq i < d$ we have*

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^{unm}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^{unm} + (x_1, \dots, x_i)}$$

Moreover if $i \neq 1$ we have

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^{unm}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^{unm}}$$

In fact, any sop $x_1, \dots, x_d$ in $\mathfrak{m}^N$ for $N \gg 0$ has such a property.

The following lemma, which we will use later, is also used to show Prop. 14.

Lemma 15 (6.9 [14]). *In the situation of Prop. 14, we have $(x_1, \dots, x_i) : x_{i+1} = (x_1, \dots, x_i)^{unm}$ for $0 \leq i < d$, by replacing the sequence $x_1, \dots, x_d$ by high powers, if necessary.*

Proof of Prop. 14. By Schenzel's formula (3.3) [12], we have

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i) : x_{i+1}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i) : x_j + (x_1, \dots, x_i)}$$

for $i = 0, \dots, d-1$. Thus we obtain the desired result by Lemma 15. Notice that for $i = 2, \dots, d-1$, we have $(x_1, \dots, x_i) \subset \sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i) : x_j$. $\square$

Notice that the equidimensional hull of a graded ideal is again graded. Then, by considering a graded version of Prop. 3.3 in [12], which Prop. 14 bases on, we easily deduce the following.

Corollary 16. *Assume that $(R, \mathfrak{m})$ is generalized Cohen-Macaulay. Then there exists a homogeneous sop $x_1, \dots, x_d$, ($d = \dim R$) such that for $0 \leq i < d$ we have*

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^{unm}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^{unm} + (x_1, \dots, x_i)} (\delta_i)$$

where $\delta_i = \sum_{j=1}^i \deg(x_j)$. Moreover if $i \neq 1$ we have

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^{unm}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^{unm}}(\delta_i).$$

Theorem 17. *Let $(R, \mathfrak{m})$ be an $\mathbb{N}$ -graded Noetherian K -algebra of dimension $d = \dim R$ and $\text{char } K = p > 0$. Assume that R is an equidimensional isolated non F -rational singularity. Then for any homogeneous sop $x_1, \dots, x_d \in \mathfrak{m}^N$ with $N \gg 0$, we have*

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{\lim}}(\delta_i) \cong \frac{(x_1^n, \dots, x_i^n)^*}{(x_1^n, \dots, x_i^n)^{\lim}}(n\delta_i)$$

for $i = 0, \dots, d-1$ and $n \in \mathbb{N}$, where we define $\delta_i = \sum_{j=1}^i \deg x_j$.

Proof. Let $x_1, \dots, x_d$ be a homogeneous sop. Since R is generalized Cohen-Macaulay, Cor. 7 allows us to assume that it is a USD-sequence by replacing it, if necessary, by higher powers. For $i = 0, \dots, d-1$, we compute

$$\begin{aligned} H_{\mathfrak{m}}^i(R) &\cong \frac{(x_1, \dots, x_i)^{unm}}{\sum_{j=1}^i (x_1, \dots, \hat{x}_j, \dots, x_i)^{unm} + (x_1, \dots, x_i)}(\delta_i) \quad \text{by Cor.16} \\ &\cong \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{germ} + (x_1, \dots, x_i)}(\delta_i) \quad \text{by Cor.12} \\ &\cong \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{\lim}}(\delta_i) \quad \text{by Prop. 13} \end{aligned}$$

The isomorphism

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1^n, \dots, x_i^n)^*}{(x_1^n, \dots, x_i^n)^{\lim}}(n\delta_i) \quad (n \in \mathbb{N})$$

is immediate since, for a USD-sequence $x_1, \dots, x_i$, its power $x_1^n, \dots, x_i^n$ ($n \in \mathbb{N}$) is also a USD-sequence. $\square$

As an immediate application of this representation, we recover some of the known results in a slightly general way.

Corollary 18 (cf. (2.7) [11]). *Let $(R, \mathfrak{m})$ be an equidimensional isolated non F -rational singularity and let $I = (x_1, \dots, x_i)$, $i < d = \dim R$, be a homogeneous parameter ideal such that with $I \subset \mathfrak{m}^\ell$ for $\ell \gg 0$. Then for an integer N_i the following are equivalent*

- (i) $[H_{\mathfrak{m}}^i(R)]_n = 0$ for all $n < N_i$.
- (ii) $I^* \subseteq I^{\lim} + R_{\geq \delta_i + N_i}$ where $\delta_i = \sum_{j=1}^i \deg(x_j)$.

In a special case, we can give a lower bound of the vanishing degree N_i . For an $\mathbb{N}$ -graded module M , we set $b(M) = \min\{i \mid M_i \neq 0\}$. If $(R, \mathfrak{m})$ is a generalized Cohen-Macaulay local ring, we have

$$[H_{\mathfrak{m}}^i(R)]_n = 0 \quad \text{for } n < -it, \quad i < \dim R,$$

if there exists a homogeneous sop $x_1, \dots, x_d$ as in Th. 17 with $t = \min\{\deg x_1, \dots, \deg x_d\}$ ((2.4)[8]). The following result gives a refinement for isolated non F -rational singularities.

Corollary 19. *Let K be an algebraically closed field of char $K = p > 0$ and let $(R, \mathfrak{m})$ be a Noetherian standard $\mathbb{N}$ -graded domain with $R_0 = K$ and $\mathfrak{m} = \bigoplus_{n>0} R_n$. Assume that R is an isolated non F -rational singularity and let*

$$t = \min\{N \in \mathbb{N} \mid \mathbf{x} = x_1, \dots, x_d \in \mathfrak{m}^N \text{ where } \mathbf{x} \text{ is as in Th. 17}\}.$$

Then for all $i < d = \dim R$ and for all $n < -(i-1)t$ we have $[H_{\mathfrak{m}}^i(R)]_n = 0$.

Proof. By Th. 17 we can find an sop $x_1, \dots, x_d \in \mathfrak{m}^N$ such that for every $i < d$

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{\lim}}(\delta_i) \quad \text{where} \quad \delta_i = \sum_{j=1}^i \deg(x_j)$$

The lower bound of the degrees of the non-zero elements in $(x_1, \dots, x_i)$ is $\min\{\deg x_1, \dots, \deg x_i\}$, so that we have $b((x_1, \dots, x_i)^*(\delta_i)) = b((x_1, \dots, x_i)^*) - \delta_i$, which is, by Exer. 4.5 and 4.6 in [9], equal to $b((x_1, \dots, x_i)) - \delta_i = \min\{\deg x_1, \dots, \deg x_i\} - \delta_i \geq -(i-1)t$ as required. $\square$

For another application, we can consider, under the condition of Theorem 17, the natural map $\rho_i : H_{\mathfrak{m}}^i(R) \rightarrow H_{\mathfrak{m}}^{i+1}(R)$ for $i = 0, \dots, d-2$ defined by $\rho_i(r + (x_1, \dots, x_i)^{\lim}) = r + (x_1, \dots, x_{i+1})^{\lim}$ with $r \in (x_1, \dots, x_i)^*$. For the degree preserving homomorphism, we can also consider $\psi_i(r + (x_1, \dots, x_i)^{\lim}) = rx_{i+1} + (x_1, \dots, x_{i+1})^{\lim}$. Unfortunately, they are all trivial.

Corollary 20. *The maps ρ_i and ψ_i are trivial.*

Proof. We first show that $(x_1, \dots, x_i) : x_{i+1} \subset (x_1, \dots, x_{i+1})^{\lim}$ for $i < d$. Take any $r + (x_1, \dots, x_i)^{\lim} \in \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{\lim}} \cong H_{\mathfrak{m}}^i(R)$. By Prop. 11 and Lemma 15, we have $r \in (x_1, \dots, x_i)^* = (x_1, \dots, x_i) : x_{i+1}$, i.e., $x_{i+1}r \in (x_1, \dots, x_i)$, so that, for any $s \in \mathbb{N}$, we have $x_1^s \cdots x_i^s x_{i+1}^s r \in (x_1^{s+1}, \dots, x_{i+1}^{s+1})$, which implies that $r \in (x_1, \dots, x_{i+1})^{\lim}$. Hence $\rho_i = 0$. $\psi_i = 0$ is similarly proved. $\square$

3. TIGHT CLOSURE OF ZEROS AND ISOLATED SINGULARITIES

In this section, we consider the question of how the tight closure of zero in the highest local cohomology $H_{\mathfrak{m}}^d(R)$ controls the vanishing of the lower local cohomologies of isolated singularities.

Definition 12 (tight closure of zero). *For a R -module M , we define*

$$(0)_M^* = \{x \in M \mid \text{there exists } c \in R^o \text{ such that } cx^q = 0 \text{ in } M \text{ for } q = p^e \gg 0\}$$

and call it tight closure of 0 in M .

We are particularly interested in the tight closure of zero in the highest local cohomology $H_{\mathfrak{m}}^d(R)$, $d = \dim R$. We give a graded version of the result by Goto and Nakamura.

Proposition 21. *Let $(R, \mathfrak{m})$ be an $\mathbb{N}$ -graded isolated non F -rational singularity. Suppose that a homogeneous sop $\mathbf{x} = x_1, \dots, x_d$ forms a USD-sequence. Then, by replacing $\mathbf{x}$ by a higher power $\mathbf{x}^N$ ($N > 0$) if necessary, we have*

$$(0)_{H_{\mathfrak{m}}^d(R)}^* = \bigcup_{n>0} Z_n(n\delta) \quad \text{where} \quad Z_n = \frac{(x_1^n, \dots, x_d^n)^*}{(x_1^n, \dots, x_d^n)^{\lim}},$$

where $\delta = \sum_{j=1}^d \deg(x_j)$.

Proof. By considering the graded version of Prop. 2.1 [4], we have

$$(0)_{H_{\mathfrak{m}}^d(R)}^* = \bigcup_{n>0} Z'_n(n\delta) \quad \text{where} \quad Z'_n = \frac{(x_1^n, \dots, x_d^n)^*}{\sum_{j=1}^n (x_1^n, \dots, \hat{x}_j^n, \dots, x_d^n) : x_j^n + (x_1^n, \dots, x_d^n)}.$$

Then by Cor. 12, Prop. 13 and Lemma 15, we obtain the desired result. $\square$

Notice that, for $i < d = \dim R$, we have a natural map

$$H_{\mathfrak{m}}^i(R) \cong \frac{(x_1, \dots, x_i)^*}{(x_1, \dots, x_i)^{\lim}}(\delta_i) \longrightarrow \frac{(x_1, \dots, x_d)^*}{(x_1, \dots, x_d)^{\lim}}(\delta_d) \subset (0)^* \subset H_{\mathfrak{m}}^d(R)$$

by sending $r + (x_1, \dots, x_i)^{\lim}$ to $r + (x_1, \dots, x_d)^{\lim}$ or as a degree preserving map $rx_{i+1} \cdots x_d + (x_1, \dots, x_d)^{\lim}$. But they are actually 0-maps, which can be proved similarly to Cor. 20.

Now we consider relation between the tight closures of zero of $H_{\mathfrak{m}}^d(R)$ and $H_{\mathfrak{m}}^{d-1}(\overline{R})$, where $\overline{R}$ is a hypersurface intersetion of R . It is well known that Flenner's Bertini theorem also holds for positive characteristics, which is, as far as the author is concerned, not stated in the literature. We give the statement for the readers convenience.

Proposition 22 (cf. Satz (4.1) [3]). *Let k be an infinite field and $(R, \mathfrak{m})$ be a Noetherian local k -algebra, whose residue class ring $K := R/\mathfrak{m}$ is separable over k . Let $x_1, \dots, x_d \in \mathfrak{m}$ be such that $I = (x_1, \dots, x_d)$ is an $\mathfrak{m}$ -primary ideal. Assume that R_P is regular for all $P \in \operatorname{Spec} R - \{\mathfrak{m}\}$. Then for a general linear form $x_\alpha = \sum_{i=1}^d \alpha_i x_i$ with $\alpha \in k^d$ and any $P \in \operatorname{Spec} R/x_\alpha - \{\mathfrak{m}'\}$ where $\mathfrak{m}' = \mathfrak{m}/x_\alpha R$, $(R/x_\alpha R)_P$ is again regular.*

Let $(R, \mathfrak{m})$ be an isolated singularity with $d = \dim R$. Then, by Prop. 5, 22 and 21 together with Cor. 7, we can choose a homogeneous sop $x_1, \dots, x_d$, which is a USD-sequence, such that $\overline{R} = R/x_d R$ is also an isolated singularity and

$$(0)_{H_{\mathfrak{m}}^d(R)}^* = \bigcup_{n>0} \frac{(x_1^n, \dots, x_d^n)^*}{(x_1^n, \dots, x_d^n)^{\lim}}(n\delta_d) \quad \text{and} \quad (0)_{H_{\mathfrak{m}}^{d-1}(\overline{R})}^* = \bigcup_{n>0} \frac{(\overline{x}_1^n, \dots, \overline{x}_{d-1}^n)^*}{(\overline{x}_1^n, \dots, \overline{x}_{d-1}^n)^{\lim}}(n\delta_{d-1})$$

where $\delta_j = \sum_{k=1}^j \deg(x_k)$ and $\overline{x}_j$ is the image of x_j in $\overline{R}$. Now we can prove

Proposition 23. *Let $(R, \mathfrak{m})$ be an $\mathbb{N}$ -graded isolated non F -rational singularity over the field K of char $K = p > 0$ with $R_0 = K$. Let $d = \dim R$. Assume that $x_1, \dots, x_d$ is a homogeneous sop and consider the natural surjection*

$$\varphi : R \longrightarrow \overline{R} = R/x_d R,$$

whose image $\varphi(r)$ will be denoted by $\overline{r}$. Then,

(i) For any $n \in \mathbb{Z}$ we have

$$\text{if } \left[\frac{(x_1, \dots, x_d)^*}{(x_1, \dots, x_d)^{\lim}} \right]_n \neq 0, \quad \text{then } \left[\frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}} \right]_n \neq 0,$$

(ii) Assume also that $x_1, \dots, x_d$ is a USD-sequence and that $R/x_d^\ell R$ is a non F -rational singularity for all $\ell \in \mathbb{N}$. If $[(0)_{H_m^d(R)}^*]_n \neq 0$ for $n \in \mathbb{Z}$, then there exists $\ell \in \mathbb{N}$ such that $[(0)_{H_m^{d-1}(R/x_d^\ell R)}^*]_{n+a} \neq 0$, where $a = \ell \cdot \deg(x_d)$.

Proof. We first show (i). By persistence of tight closure (cf. Th. 2.3 [9]) we have $\varphi((x_1, \dots, x_d)^*) \subset (\bar{x}_1, \dots, \bar{x}_d)^* = (\bar{x}_1, \dots, \bar{x}_{d-1})^*$. Thus φ induces the homomorphism

$$\bar{\varphi} : (x_1, \dots, x_d)^* \longrightarrow \frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}}.$$

We now show $\text{Ker } \bar{\varphi} \subset (x_1, \dots, x_d)^{\lim}$. Let $r \in \text{Ker } \bar{\varphi}$ be arbitrary. Then $\bar{r} := \varphi(r) \in (\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}$ and we have $\bar{x}_1^s \cdots \bar{x}_{d-1}^s \cdot \bar{r} \in (\bar{x}_1^{s+1}, \dots, \bar{x}_{d-1}^{s+1})$ for some $s \in \mathbb{N} \cup \{0\}$, which implies that $x_1^s \cdots x_{d-1}^s r \in (x_1^{s+1}, \dots, x_{d-1}^{s+1}, x_d)$ and then $x_1^s \cdots x_d^s r \in (x_1^{s+1}, \dots, x_d^{s+1})$. Hence $r \in (x_1, \dots, x_d)^{\lim}$ as required. Consequently, we have the following diagram:

$$(1) \quad \frac{(x_1, \dots, x_d)^*}{(x_1, \dots, x_d)^{\lim}} \xleftarrow{\text{nat}} \frac{(x_1, \dots, x_d)^*}{\text{Ker } \bar{\varphi}} \xrightarrow{\hat{\varphi}} \frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}}$$

where nat is the natural surjection and $\hat{\varphi}$ is the induced embedding. Taking the degree n fraction of this diagram, we immediately know that (i) holds.

Now we show (ii). Assume that $[(0)_{H_m^d(R)}^*]_n \neq 0$. Then by Prop. 21 there exists $\ell \in \mathbb{N}$ such that

$$0 \neq \left[\frac{(x_1^\ell, \dots, x_d^\ell)^*}{(x_1^\ell, \dots, x_d^\ell)^{\lim}} (\ell \cdot \delta_d) \right]_n = \left[\frac{(x_1^\ell, \dots, x_d^{\ell l})^*}{(x_1^\ell, \dots, x_d^\ell)^{\lim}} \right]_{n+\ell \cdot \delta_d}.$$

Thus by (i) with $x_1, \dots, x_d$ replaced by $x_1^\ell, \dots, x_d^\ell$, we have

$$0 \neq \left[\frac{(\bar{x}_1^\ell, \dots, \bar{x}_{d-1}^\ell)^*}{(\bar{x}_1^\ell, \dots, \bar{x}_{d-1}^\ell)^{\lim}} \right]_{n+\ell \cdot \delta_d} = \left[\frac{(\bar{x}_1^\ell, \dots, \bar{x}_{d-1}^\ell)^*}{(\bar{x}_1^\ell, \dots, \bar{x}_{d-1}^\ell)^{\lim}} (\ell \cdot \delta_{d-1}) \right]_{n+\ell \cdot \deg(x_d)}$$

Then we obtain the required result by Prop. 21. $\square$

Prop. 23 assumes that hypersurface intersections of isolated non F -rational singularities are again isolated non F -rational singularities. However, F -rationality behaves rather badly with hypersurface intersections, since an F -rational local ring has negative a -invariant (see [2]) but hypersurface intersection increases a -invariant. But for isolated singularity, which is a special case of isolated non F -rational singularity, hypersurface intersections behave well thanks to Bertini type theorems.

Now we come to the main result of this section.

Theorem 24. *Let $(R, \mathfrak{m})$ be a reduced $\mathbb{N}$ -graded equidimensional isolated singularity over the field K of char $K = p > 0$ with $R_0 = K$, $d = \dim R$ and $\sharp K = \infty$. Assume that $[(0)_{H_m^d(R)}^*]_n \neq 0$ for some $n \in \mathbb{Z}$ and consider a homogeneous regular element x , $a = \deg(x)$, that is also a part of a USD-sequence. Then*

- (i) If the multiplication $[H_{\mathfrak{m}}^d(R)]_n \xrightarrow{x} [H_{\mathfrak{m}}^d(R)]_{n+a}$ is injective, then we have $[H_{\mathfrak{m}}^{d-1}(R)]_{n+\ell \cdot a} \neq 0$ for some $\ell \in \mathbb{N}$.
- (1) Otherwise, if there is a USD sequence $x_1, \dots, x_d$ with $x = x_d$ such that the degree $n + \delta_d$ fragment of $(x_1, \dots, x_d)^{\lim} - \varphi^{-1}((\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim})$ is nonempty, where $\varphi : R \rightarrow R/x_d R$ is the natural surjection whose image of $r \in R$ is denoted by $\bar{r}$. Then we have $[H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \neq 0$.

Proof. Let $x_1, \dots, x_d \in \mathfrak{m}$ be a USD-sequence, whose existence is assured by Prop. 6. Since R is reduced, we have $\text{depth } R > 0$ and thus we can take x_d general enough to be R -regular. Consider the short exact sequence:

$$0 \longrightarrow R(-a) \xrightarrow{x_d} R \xrightarrow{\varphi} \bar{R} \longrightarrow 0, \quad a := \deg(x_d)$$

where we set $\bar{R} := R/(x_d)R$ and $\varphi : R \rightarrow R/(x_d)R$ is the natural map. Also by Prop. 22, we can assume that $\bar{R}$ is again an isolated singularity.

Since x_d is a part of the USD-sequence, it is standard by Prop. 4, so that we have $x_d H_{\mathfrak{m}}^{d-1}(R) = 0$. Thus, from the above short exact sequence, we obtain the long exact sequence

$$0 \longrightarrow H_{\mathfrak{m}}^{d-1}(R) \longrightarrow H_{\mathfrak{m}}^{d-1}(\bar{R}) \xrightarrow{\psi} H_{\mathfrak{m}}^d(R)(-a) \xrightarrow{x_d} H_{\mathfrak{m}}^d(R) \longrightarrow 0,$$

and we consider the degree $n + a$ fragment:

$$0 \longrightarrow [H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \longrightarrow [H_{\mathfrak{m}}^{d-1}(\bar{R})]_{n+a} \xrightarrow{\psi} [H_{\mathfrak{m}}^d(R)]_n \xrightarrow{x_d} [H_{\mathfrak{m}}^d(R)]_{n+a} \longrightarrow 0.$$

Now we show (i). Assume that the multiplication by x_d is injective, we have an isomorphism $[H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \cong [H_{\mathfrak{m}}^{d-1}(\bar{R})]_{n+a}$. By Prop. 21 we have the inclusion

$$\frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}}(\delta_{d-1}) \subset (0)_{H_{\mathfrak{m}}^{d-1}(\bar{R})}^* \subset H_{\mathfrak{m}}^{d-1}(\bar{R})$$

so that, if

$$\left[\frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}}(\delta_{d-1}) \right]_{n+a} \neq 0,$$

then we have $[H_{\mathfrak{m}}^{d-1}(\bar{R})]_{n+a} \cong [H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \neq 0$. Then (i) follows by Prop. 23(ii) by replacing, if necessary, x_d by x_d^ℓ for some $\ell \in \mathbb{N}$.

Next we show (ii). If $x_d \in \mathfrak{m}^N$ for $N \gg 0$, x_d is a parameter test element by Prop. 3. By Prop 4.4 [13] we know $J \subset \text{Ann}_R((0)_{H_{\mathfrak{m}}^d(R)}^*)$ where J is the parameter test ideal. Thus $x_d : [H_{\mathfrak{m}}^d(R)]_n \xrightarrow{x_d} [H_{\mathfrak{m}}^d(R)]_{n+a}$ is not injective if $[(0)_{H_{\mathfrak{m}}^d(R)}^*]_n \neq 0$. Thus we have the exact sequence

$$0 \longrightarrow [H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \longrightarrow [H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \xrightarrow{\psi} \text{Im } \psi \longrightarrow 0$$

where $0 \neq [(0)_{H_{\mathfrak{m}}^d(R)}^*]_n \subset \text{Im } \psi$. By considering the Čech complex representation of local cohomology, we know that, for

$$\eta = \left[\frac{\bar{r}}{\bar{x}_1 \cdots \bar{x}_{d-1}} \right] \in H_{\mathfrak{m}}^{d-1}(\bar{R})$$

we have

$$\psi(\eta) = \left[\frac{r}{x_1 \cdots x_d} \right] \in H_{\mathfrak{m}}^d(R).$$

Also we know that

$$\begin{aligned} \text{Ker } \psi &= \left\{ \eta = \left[\frac{\bar{r}}{\bar{x}_1 \cdots \bar{x}_{d-1}} \right] \mid \left[\frac{r}{x_1 \cdots x_d} \right] = 0 \text{ in } H_{\mathfrak{m}}^d(R) \right\} \\ &= \left\{ \eta = \left[\frac{\bar{r}}{\bar{x}_1 \cdots \bar{x}_{d-1}} \right] \mid r \in (x_1, \dots, x_d)^{\lim} \right\} \end{aligned}$$

by Prop. 9. We want to show that $\text{Ker } \psi \neq 0$. Now consider the diagram (1) in the proof of Prop. 23. We consider the submodule

$$L := \text{Im } \hat{\varphi} \subset \frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}} \subset (0)_{H_{\mathfrak{m}}^{d-1}(\bar{R})}^*(-\delta_{d-1}).$$

Then $\psi([L]_{n+\delta_d}) \subset [(0)_{H_{\mathfrak{m}}^d(R)}^*]_n$, since

$$\psi(\eta) = \left[\frac{r}{x_1 \cdots x_d} \right] \in (0)_{H_{\mathfrak{m}}^d(R)}^*$$

if and only if $r \in (x_1, \dots, x_d)^*$ (see for example [10]). Now, if $r \in (x_1, \dots, x_d)^{\lim} - \text{Ker } \bar{\varphi}$, then the image of r in $(x_1, \dots, x_d)^*/\text{Ker } \bar{\varphi}$ in the diagram (1) give the non-zero element $\eta \in (\bar{x}_1, \dots, \bar{x}_{d-1})^*/(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}$, which contributes to a non-zero element in $H_{\mathfrak{m}}^{d-1}(R)$. Moreover we have $\psi(\eta) = 0$, since $\psi(\eta)$ is the image of r in $(x_1, \dots, x_d)^*/(x_1, \dots, x_d)^{\lim}$. Thus $\text{Ker } \psi \neq 0$ so that we must have $[H_{\mathfrak{m}}^{d-1}(R)]_{n+a} \neq 0$ as required. $\square$

Remark 1. As we saw in the proof, the condition (i) of Th. 24 is satisfied only when $\deg(x)$ is small large enough. On the other hand, Prop. 6 tells that, generally speaking, in order to obtain a part of USD-sequence we need to choose elements of high degree. Thus (i) could be satisfied in relatively few cases.

Also, since $[H_{\mathfrak{m}}^{d-1}R]_k = 0$ for $k \gg 0$, Th. 24 means that $(x_1, \dots, x_d)^{\lim}$ is the preimage of $(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}$ via the natural map φ when $\deg(x_d)$ is large enough.

The following result describes the injectivity of the morphism *nat* in the diagram (1).

Corollary 25. *Let $(R, \mathfrak{m})$ be a reduced $\mathbb{N}$ -graded equidimensional isolated singularity over the field K of char $K = p > 0$ with $R_0 = K$, $d = \dim R$ and $\sharp K = \infty$. For an sop $x_1, \dots, x_d \in \mathfrak{m}^N$ with $N \gg 0$ and for every $n \in \mathbb{Z}$ with $[(0)_{H_{\mathfrak{m}}^d(R)}^*]_n \neq 0$, we have*

$$\left[\frac{(x_1, \dots, x_d)^*}{(x_1, \dots, x_d)^{\lim}} \right]_{n+\delta_d} \subset \left[\frac{(\bar{x}_1, \dots, \bar{x}_{d-1})^*}{(\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim}} \right]_{n+\delta_d}$$

where $\bar{x}_j$ is the image of x_j in $\bar{R} := R/x_d R$.

Proof. Since $[H_{\mathfrak{m}}^{d-1}(R)]_{n+a} = 0$ for $a \gg 0$, we know that the conditions (i) and (ii) in Th. 24 do not hold for sop's with high enough degrees, which implies that $[(x_1, \dots, x_d)^{\lim}]_{n+\delta_d} = [\varphi^{-1}((\bar{x}_1, \dots, \bar{x}_{d-1})^{\lim})]_{n+\delta_d}$ as required. $\square$

REFERENCES

- [1] P. Deligne, Pierre and L. Illusie, Luc. Relèvements modulo p^2 et décomposition du complexe de de Rham. *Invent. Math.* 89 (1987), no. 2, 247–270. MR0894379
- [2] R. Fedder and K.-i. Watanabe, A characterization of F -regularity in terms of F -purity. *Commutative algebra* (Berkeley, CA, 1987), 227–245, Math. Sci. Res. Inst. Publ., 15, Springer, New York, 1989. MR1015520
- [3] H. Flenner, Die Sätze von Bertini für lokale Ringe. *Math. Ann.* 229 (1977), no. 2, 97–111. MR0460317
- [4] S. Goto and Y. Nakamura, The bound of the difference between parameter ideals and their tight closures. *Tokyo J. Math.* 25 (2002), no. 1, 41–48. MR1908212
- [5] S. Goto and K. Yamagishi, The theory of unconditioned strong d -sequences and modules of finite local cohomology, preprint
- [6] N. Hara, A characterization of rational singularities in terms of injectivity of Frobenius maps. *Amer. J. Math.* 120 (1998), no. 5, 981–996. MR1646049
- [7] H. Hara and K.-i. Watanabe, The injectivity of Frobenius acting on cohomology and local cohomology modules. *Manuscripta Math.* 90 (1996), no. 3, 301–315. MR1397659
- [8] L. T. Hoa and C. Miyazaki, Bounds on Castelnuovo-Mumford regularity for generalized Cohen-Macaulay graded rings. *Math. Ann.* 301 (1995), no. 3, 587–598. MR1324528
- [9] C. Huneke, *Tight Closure and Its Applications*, CBMS Regional Conference Series in Mathematics, 88. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1996. x+137 pp. ISBN: 0-8218-0412-X MR1377268
- [10] C. Huneke, Tight closure, parameter ideals, and geometry. *Six lectures on commutative algebra* (Bellaterra, 1996), 187–239, Progr. Math., 166, Birkhäuser, Basel, 1998. MR1648666
- [11] C. Huneke and K. Smith, Tight closure and the Kodaira vanishing theorem. *J. Reine Angew. Math.* 484 (1997), 127–152. MR1437301
- [12] P. Schenzel, Standard systems of parameters and their blowing-up rings. *J. Reine Angew. Math.* 344 (1983), 201–220. MR0716256
- [13] K. Smith, Test ideals in local rings. *Trans. Amer. Math. Soc.* 347 (1995), no. 9, 3453–3472. MR1311917
- [14] K. Smith, Vanishing, singularities and effective bounds via prime characteristic local algebra. *Algebraic geometry—Santa Cruz 1995*, 289–325, Proc. Sympos. Pure Math., 62, Part 1, Amer. Math. Soc., Providence, RI, 1997. MR1492526
- [15] M. Raynaud, Contre-exemple au “vanishing theorem” en caractéristique $p > 0$. C. P. Ramanujam—a tribute, pp. 273–278, Tata Inst. Fund. Res. Studies in Math., 8, Springer, Berlin-New York, 1978. MR0541027
- [16] J. Velez, Openness of the F -rational locus and smooth base change. *J. Algebra* 172 (1995), no. 2, 425–453. MR1322412

YUKIHIRO TAKAYAMA, DEPARTMENT OF MATHEMATICAL SCIENCES, RITSUMEIKAN UNIVERSITY, 1-1-1 NOJIHIGASHI, KUSATSU, SHIGA 525-8577, JAPAN
E-mail address: takayama@se.ritsumei.ac.jp